

White Coat Companion 2021 Errata

Updated: 12/15/21

Page Number	Type (Date Added)	Location	Details																
Cardiology																			
Cards 27	Update (12/15/21)	Carotid Artery Disease	<p>Treatment guidelines updated for both symptomatic and asymptomatic disease.</p> <table border="1"> <thead> <tr> <th>Symptomatic</th> <th>Intervention</th> </tr> </thead> <tbody> <tr> <td>100% Stenosis</td> <td>- Medical management</td> </tr> <tr> <td>70-99% Stenosis</td> <td>- Carotid endarterectomy (ideally performed between 3-14 days after stroke symptoms) - Carotid artery stenting (if high risk for surgery or lesion is not surgically accessible)</td> </tr> <tr> <td>50-69% Stenosis</td> <td>- Carotid endarterectomy (if reasonable life expectancy)</td> </tr> <tr> <td>< 50% Stenosis</td> <td>- Medical management</td> </tr> <tr> <th>Asymptomatic</th> <th>Intervention</th> </tr> <tr> <td>≥ 70% Stenosis</td> <td>- Carotid endarterectomy (if low surgical risk/long life expectancy) or intensive medical therapy</td> </tr> <tr> <td>< 70%</td> <td>- Medical management (statin, antiplatelet, BP control, etc)</td> </tr> </tbody> </table>	Symptomatic	Intervention	100% Stenosis	- Medical management	70-99% Stenosis	- Carotid endarterectomy (ideally performed between 3-14 days after stroke symptoms) - Carotid artery stenting (if high risk for surgery or lesion is not surgically accessible)	50-69% Stenosis	- Carotid endarterectomy (if reasonable life expectancy)	< 50% Stenosis	- Medical management	Asymptomatic	Intervention	≥ 70% Stenosis	- Carotid endarterectomy (if low surgical risk/long life expectancy) or intensive medical therapy	< 70%	- Medical management (statin, antiplatelet, BP control, etc)
Symptomatic	Intervention																		
100% Stenosis	- Medical management																		
70-99% Stenosis	- Carotid endarterectomy (ideally performed between 3-14 days after stroke symptoms) - Carotid artery stenting (if high risk for surgery or lesion is not surgically accessible)																		
50-69% Stenosis	- Carotid endarterectomy (if reasonable life expectancy)																		
< 50% Stenosis	- Medical management																		
Asymptomatic	Intervention																		
≥ 70% Stenosis	- Carotid endarterectomy (if low surgical risk/long life expectancy) or intensive medical therapy																		
< 70%	- Medical management (statin, antiplatelet, BP control, etc)																		
Pulm/Crit																			
Pulm 18	Clarification (3/8/21)	Acute Respiratory failure	Non-rebreather can go up to 15 L/min flow																
Pulm 19	Minor error (3/8/21)	Mechanical Ventilation	Units on PEEP is cmH2O																
Renal																			
Endocrine																			
Endo 34	Error (6/3/21)	Metformin	INCREASES peripheral glucose uptake																
Endo 35	Clarification (6/28/21)	GLP-1	Are mainly delivered subQ (not oral), although a new oral GLP-1 agonist was just approved.																
GI																			
GI 26	Clarification (6/3/21)	External Hemorrhoids	Can bleed and cause minor irritation. Severe pain is generally caused by thrombosed external hemorrhoids.																
Heme-Onc																			

Heme 13	Minor error (6/3/21)	TTP	ADAMTS13 (typo in spelling)
Rheum/MSK			
Rheum 11	Error (6/3/21)	Myasthenia	Anti AChR antibodies (not anti BM)
Rheum 12	Error (6/4/21)	PM/DM	Under diagnosis, classic skin rash is associated with dermatomyositis
Derm			
ID			
Neurology			
Neuro 10	Minor error (12/15/21)	TBI	Severe TBI GCS < 9
Neuro 25	Update (12/15/21)	LEMS	3, 4 diaminopyridine is first line for treatment of LEMS
Primary Care/EM			
PC/EM 1	Update (3/10/21)	Adult Screening	Low dose CT scan for: Adults aged 50 to 80 years who have a 20 pack-year smoking history and currently smoke or have quit within the past 15 years (updated 2021 USPSTF guidelines)
Psychiatry			
Pediatrics			
Surgery			
OB-GYN			
OBGYN 3	Update (6/3/21)	Prenatal Testing	Pap smears should be performed at the same routine intervals that are recommended for non-pregnant women
OBGYN 17	Error (10/7/21)	Choriocarcinoma	Proliferation of cytotrophoblast and syncytiotrophoblast with OUT chorionic villi
OB-GYN 64	Update (6/3/21)	AUB	Infrequent bleeding is less frequently than every 38 days (not 28)

